

## Medication List

If you are using any of the medications listed below, their use should be discontinued a minimum of two weeks before surgery. You can resume taking them 5 days after surgery. If your doctor ordered any of these medications, consult him or her before you stop.

Advil (1)	Dasin	Nalfon (1)
Alka Seltzer	Diflunisal (3)	Naprosyn (1)
Anacin	Dipyridamole (1)	Naproxen (3)
Anaprox (3)	Dolcin	Nuprin (1)
Anexsia with Codeine	Dolene Compound	Oxalid (21)
Anturane (2)	Dolobid (3)	Oxyphenbutazone
APC with Codeine	Double – A	Percodan
Arthritis Bayer	Duradyne	Persantine
Arthritis Pain Formula	Dynosal	Phenybutazone
Ascriptin	Easprin	Piroxicam (14)
Asper Buf	Ecotrin	Presalin
Aspercin	Emagrin	Rufen (1)
Aspergum	Emprin	Salagen
Aspirin	Equagesic	Saleto
Axotal	Equazine – M	Salocol
Azolid (21)	Excedrin	Sine-Off
Bayer	Feldene	SK-65 Compound
BC Tabs or Powder	Fenoprofen (1)	Speedrin
Buf Tab	Fiorinal	Sulfinpyrazone
Buff A or Compound	Gemnisyn	Sulindac (3)
Buffaprin	Goodys Headache Pwdr	Supac
Bufferin	Ibuprofen (1)	Synalgos
Bufferin EX-Strength	Indameth (1)	Talwin Compound
Buffets II	Indocin (1)	Tandearil (21)
Buffex	Indocin SR (1)	Tolectin (1)
Butagen (21)	Indomed (1)	Tolmetin (1)
Butazolidin (21)	Indo – Lemmon (1)	Trigesic
Butazone (21)	Indomethacin (1)	Vanquish
Cama	Lanorinol	Zorpin
Capron	Magnaprin	
Clinoril (3)	Marnal	
Codoxy	Measurin	
Congesprin	Mefanamic Acid (1)	
Cope	Mepro-Analgesic #2	
Coricidin	Mepro Compound	
Cosprin	Micrainin	
CP-2	Midol Max-Strength	
Darvon Compound	Momentum	
Darvon with ASA	Motrin (1)	

If you have any questions or concerns regarding medications listed, call your doctor.